arXiv:physics/0006068v1 [physics.space-ph] 27 Jun 2000

ENHANCED PITCH ANGLE DIFFUSION DUE TO ELECTRON-WHISTLER INTERACTIONS DURING DISTURBED TIMES

W. Wykes, S. C. Chapman, and G. Rowlands

Space and Astrophysics Group, University of Warwick, Coventry, UK

ABSTRACT

During periods of increased magnetospheric activity, whistler wave emissions have been observed with increased wave amplitudes. We consider a pitch angle diffusion mechanism that is shown to scale with whistler wave amplitude and hence is 'switched on' during these periods of intense activity.

We consider the interaction between relativistic electrons and two oppositely directed, parallel propagating whistler mode waves. We show that for intense whistlers this process is stochastic and results in strong electron pitch angle diffusion.

We show that the interaction is rapid, occur on timescales of the order of tens of electron gyroperiods and that the interaction is sensitive to wave amplitude, wave frequency and electron energy.

Key words: Electron, Whistler, Substorms, Diffusion.

1. INTRODUCTION

Electron whistler interactions have long been considered as a mechanism for pitch angle scattering. Gyroresonance processes with near-parallel propagating waves (e.g. Kennel and Petschek (1966), Lyons and Williams (1984)) have been shown to produce pitch angle diffusion for electrons that are at resonance with a series of whistler waves (see Gendrin (1981)). We generalize resonant diffusion to include all phase space dynamics, i.e. as well as considering the resonant diffusion of trapped electrons we consider diffusion of untrapped electrons (we refer to this process as 'off-resonance' diffusion). Therefore we maximize the area in phase space contributing to pitch angle diffusion.

The underlying dynamics of the interaction between electrons and a single whistler are inherently simple, as demonstrated by Laird (1972) who derived the Hamiltonian for relativistic electrons interacting with a whistler with a single wave number. However, for a single monochromatic whistler the process is not stochastic. We introduce stochasticity by including an identical, oppositely directed whistler mode wave.

We initially consider a simplified system consisting of monochromatic whistlers in order to understand the underlying behaviour. This treatment is then extended to consider whistler wave packets, i.e. a group of waves with a range of frequencies.

We derive approximate equations in the limit of low whistler wave amplitudes and consider the degree of pitch angle diffusion for waves of different frequencies and bandwidths and for electrons of different energies.

2. EQUATIONS OF MOTION

We derive full relativistic equations of motion and approximate them in the limit of low wave amplitudes, for the case of monochromatic whistlers and wave packets. We normalize time to the electron gyrofrequency $\Omega_e = eB_0/m_e$, (where B_0 is the background magnetic field), the wave amplitude is normalized to the background magnetic field, wave frequency is normalized to the gyrofrequency and we normalize the electron velocity to the phase velocity of the waves, given by the electron dispersion relation (ignoring ion effects):

$$\frac{k^2c^2}{\omega^2} = 1 - \frac{\omega_{pe}^2}{\omega(\omega - \Omega_e)} \tag{1}$$

where ω_{pe} is the plasma oscillation frequency. Electrons can become trapped by either of the two whistlers if they are traveling close to the resonance velocity, given by:

$$\omega - \mathbf{k} \cdot \mathbf{v_r} = n\Omega_e/\gamma \tag{2}$$

where n is an integer and $\gamma = (1 - v^2/c^2)^{-1/2}$ is the relativistic factor.



Figure 1. Stroboscopic surface of section plots for the monochromatic whistler interaction. For quiet time wave amplitudes, panel a), all trajectories are regular, with resonances given by the resonance condition Equation (2). For high wave amplitudes observed during disturbed times, panels b), phase space is dominated by stochastic trajectories with regular trajectories confined to close to the resonances. The stochastic region is bounded above and below by the first regular, untrapped, trajectories away from resonance, therefore there is a limit on the diffusion of electrons in phase space.

2.1. Monochromatic Whistlers

In the limit of low wave amplitudes the full equations of motion can be reduced to:

$$\frac{d^2x}{dt^2} = \frac{2bv_{\perp 0}}{\gamma_0}\sin\left[(1/\gamma_0-\omega)t\right]\cos\left[kx\right] \quad (3)$$

where x is the distance along the background magnetic field, b is the normalized wave amplitude, $v_{\perp 0}$ is the initial perpendicular velocity and $\gamma_0 = 1/\sqrt{(1-v_{\perp 0}^2/c^2)}$.

2.2. Wave Packet Approximation

Instead of a pair of waves it is more realistic to consider the interaction of a wave packet, ie a small group of waves with a range of wave frequencies and wave numbers. We assume the wave amplitude is non-zero over wave frequency range $\Delta\omega$:

$$b(\omega) = \begin{cases} 0 & : \quad \omega < \omega_0 - \Delta\omega/2 \\ b & : \quad \omega_0 - \Delta\omega/2 < \omega < \omega_0 + \Delta\omega/2 \\ 0 & : \quad \omega > \omega_0 + \Delta\omega \end{cases} \quad (4)$$

where ω_0 is the central wave frequency of the wave packet. Integrating the monochromatic ehistler equation (3) over the frequency range $\Delta\omega$ gives the following wave packet equation:

$$\begin{aligned}\frac{d^2x}{dt^2} &= \Omega_e\frac{2bv_{\perp 0}}{\gamma_0}\sin\left[\left(\frac{1}{\gamma_0}-\omega\right)t+k_0x\right]\frac{\sin[(t-\beta x)\Delta\omega/2]}{(t-\beta x)} \\ &+ \Omega_e\frac{2bv_{\perp 0}}{\gamma_0}\sin\left[\left(\frac{1}{\gamma_0}-\omega\right)t-k_0x\right]\frac{\sin[(t+\beta x)\Delta\omega/2]}{(t+\beta x)} \quad (5)\end{aligned}$$

where $1/\beta = d\omega/dk$ is the group velocity of the waves. The wave packet equation (5) yields the monochromatic whistler equation (3) in the limit $\Delta\omega \to 0$, with amplitude $b' = b\Omega_e\Delta\omega$.

3. NUMERICAL RESULTS

The monochromatic and wave packet equations were solved numerically using a variable order, variable stepsize differential equation integrator. We consider physical parameters for the terrestrial magnetosphere at $L = 6$: gyrofrequency, $\Omega_e = 25.3\ kHz$, plasma frequency, $\omega_{pe} = 184\ kHz$, background magnetic field, $B_0 = 144\ nT$ and wave amplitude $B_\omega = 0.5\ pT$, giving a normalized wave amplitude, consistent with quiet times in the terrestrial magnetosphere, $b = 10^{-5}$ (see for example Nagano *et al.* (1996), Parrot (1994) and Summers and Ma (2000)).

The phase plots in Figure 1 are comprised of stroboscopic surfaces of section (Benettin *et al.*, 1976) to sample the full electron phase space. The initial parallel velocity was varied over the range $[-v_r, v_r]$, where v_r is the resonance velocity, given by the resonance condition, (2), for $n = 1$. All electrons were given a constant initial perpendicular velocity, with $v_\perp \approx 20v_r$ as it was found that a high velocity anisotropy was required for stochasticity.

In Figure 1 we plot parallel velocity $v_\| = dx/dt$ against phase angle ψ, where ψ is the angle between whistler propagating in a positive direction along the background field and the perpendicular velocity, $v_\perp$. In panel a) we consider a whistler wave amplitude consistent with quiet times ($b = 10^{-5}$) all trajectoriesin phase space are regular. There is little change in $v_\|$ and hence only weak pitch angle diffusion. As the wave amplitude is increased, stochastic trajectories are introduced, as the regular trajectories between the two resonances are progressively broken down. In panel b) we consider the case of intense whistler wave activity during substorms ($b = 10^{-3}$, see for example Parrot (1994) and Nagano *et al.* (1996)). The stochastic region grows to encompass the resonances as the wave amplitude is increased. Regular trajectories are confined to KAM surfaces (near-integrable trajectories with an approximate constant of the motion (Tabor, 1989)). The stochastic region is bounded by the first untrapped (regular) trajectories away from the resonances, thus there is a limit on diffusion in phase space.

As well as resonant diffusion of trapped electrons, there is diffusion of untrapped electrons throughout the stochastic region of phase space. Since, for sufficient wave amplitudes, the stochastic region can encompass the resonances, the diffusion of untrapped electrons, which we refer to as 'off-resonance' diffusion, may be enhanced over resonant diffusion. In addition we achieve pitch angle diffusion from a larger area of phase space.

Due to the time dependent nature of the wave packet equation (5) it is not possible to construct phase diagrams as in Figure 1 for the monochromatic whistler case. Instead we can consider the dynamics of single electrons. In Figure 2 we show a single trajectory solution of the wave packet equation (5), for quiet time wave amplitudes and wide ($\Delta\omega = \Omega_e/50 = 500\ Hz$) and narrow ($\Delta\omega = \Omega_e/500 = 50\ Hz$) whistler wave packets (see for example Carpenter and Sulic (1988)). We consider the change in pitch angle from an initial pitch angle of 90°. For narrow wave packets there is little change in pitch angle and the trajectory is regular. For wide wave packets the trajectory is stochastic with a large change in pitch angle ($\Delta\alpha \sim 25°$) occuring within a few tens of electron gyroperiods ($\sim 10\ ms$). We can now achieve strong pitch angle diffusion for wave amplitudes consistent with the quiet time magnetosphere.

4. PITCH ANGLE SCATTERING

Using the wave packet equation (5) with quiet time whistlers ($b = 10^{-5}$) and a relatively wide band whistler ($\Delta\omega = \Omega_e/50 = 500\ Hz$, (Carpenter and Sulic, 1988)), we can estimate the degree of pitch angle scattering. In Figures 3 and 4 we estimate the



Figure 2. *Change in pitch angle from an initial pitch angle of* 90°*, for quiet time wave amplitudes* ($b = 10^{-5}$) *and narrow and wide whistler wave packets. For narrow wave packets there is little change in pitch angle. For wide wave packets there is a large change in pitch angle* ($\Delta\alpha \sim 25°$) *occuring within a few tens of electron gyroperiods* ($\sim 10\ ms$)*, hence the interaction is rapid. Changes in pitch angle attenuate with time and the pitch angle reaches a constant value.*



Figure 3. *Log change in pitch angle,* ($\log_{10}|1+\Delta\alpha|$)*, as a function of wave frequency,* ω*, and initial pitch angle,* α_0*, for 10 keV electrons* ($v = 0.2c$) *and quiet time (low amplitude),wide band whistler wave packets* ($\Delta\omega = \Omega_e/50$)*. For high to moderate initial pitch angles* ($\alpha_0 = 50° - 90°$) *there is a change in pitch angle of up to* 40°*. For low pitch angles* ($\alpha_0 = 5° - 10°$) *the change in pitch angle is of the order of a few degrees.*

log change in pitch angle, ($\log_{10}|1+\Delta\alpha|$), as a function of wave frequency, ω, and initial pitch angle, α_0. We consider the interaction between $10keV$ electrons in Figure 3 ($100keV$ electrons in Figure 4) and wide band whistlers ($\Delta\omega = \omega_e/50$). For high to moderate initial pitch angles ($\alpha_0 = 50° - 90°$) there is a change in pitch angle of up to 40°. For low pitch angles ($\alpha_0 = 5° - 10°$) the change in pitch angle is of the order of a few degrees. In Figure 4 we see a similar degree of diffusion except that lower frequency whistler wave packets are required.



Figure 4. As for Figure 3 except for 100 keV electrons ($v = 0.5c$). Pitch angle scattering from a few degrees up to 40° *occurs although lower frequency whistler wave packets are required.*

5. DISCUSSION

We have considered electron-whistler wave particle interactions to investigate diffusion over all phase space, to include both resonant and 'off-resonance' diffusion. We have considered a simplified interaction with monochromatic whistler wave to understand the underlying behaviour and have shown that the presence of the second whistler wave introduces stochastic effects into the system. For wave amplitudes consistent with disturbed times we have shown that 'off-resonance' diffusion occurs and that resonant diffusion is unchanged.

We have considered a more realistic case of whistler wave packets and have shown that for relatively wide band whistler wave packets strong pitch angle diffusion occurs for wave amplitudes consistent with quiet, undisturbed, times. For high initial pitch angles we estimate a change in pitch angle of up to 40°, while for low pitch angles a change of a few degrees is estimated.

The effectiveness in scattering electrons of different energies is dependent on the wave frequency. Electrons with low energies (10 keV) are readily scattered by waves of around half the electron gyrofrequency, while electrons at higher energies (100 keV) are scattered by lower frequency wave ($\omega \sim \Omega_e/10$). MeV electrons would require extremely low frequency waves for efficient scattering, hence our mechanism is most efficient for electrons in the $10 - 100$ keV range.

ACKNOWLEDGEMENTS

The authors would like to acknowledge PPARC for the funding of this work.

REFERENCES

Benettin G., L. Galgani, J. M. Strelcyn. Kolmogorov entropy and numerical experiments. *Physica A, 14, 2338, 1976.*

Carpenter D. L., and D. M. Sulic. Ducted whistler propagation outside the plasmapause. *J. Geophys. Res., 93, 9731, 1988.*

Gendrin R. General Relations between wave amplification and particle diffusion in a magnetoplasma. *Rev. Geophys., 19, 171, 1981.*

Kennel C. F. and H. E. Petschek. Limit on stably trapped particle fluxes. *J. Geophys. Res., 71, 1, 1966.*

Laird M. J. Cyclotron resonance in an inhomogeneous plamsa. *J. Plasma Physics, 8, 255, 1972.*

Lyon L. R. and Williams, D. J. Quantitative Aspects of Magnetospheric Physics. *D. Reidel, Hingham, Mass., 1984.*

Nagano I., S. Yagitani, H. Kojima, H. Matsumoto. Analysis of Wave Normal and Poynting Vectors of the Chorus Emissions Observed by GEOTAIL. *J. Geomag. Geoelectr., 48, 299, 1996*

Parrot M. and C. A. Gaye. A statistical survey of ELF waves in geostationary orbit. *GRL 21, 2463, 1994.*

Summers D. and C. Ma. A model for generating relativistic electrons in the Earth's inner magnetosphere based on gyroresonant wave-particle interactions. *J. Geophys. Res., 105, 2625, 2000.*

Tabor M. Chaos and integrability in nonlinear dynamics - an introduction. *New York, Chichester: Wiley, 1989.*